New Solutions of the $2+1$ Dimensional Einstein Gravity Coupled to Maxwell Power type Non Linear Electric field with Dilaton field

Masashi Kuniyasu*
Graduate School of Science and Technology, Yamaguchi University,
Yamaguchi-shi, Yamaguchi 753–8512, Japan

January 28, 2015

arXiv:1409.4854v4 [gr-qc] 27 Jan 2015

Abstract

New solutions are derived in the $2+1$ gravity which is coupled to $|\mathcal{F}|^k$ type non-linear electric field in Maxwell Power theory with dilaton field. We obtain consistent solutions in general k case. We also investigate the behavior of the metric function with the space-time singularity. Then, we found some black hole solutions when the space-time has a singular point at $r = 0$.

1 Introduction

Recently, one of the most standard theory of gravitation is the Einstein's general theory of relativity (GR). However, GR is the classical theory, so we have to quantize the theory. To construct the quantum GR, many modified theories of gravitation had suggested. As there is a Schwarzschild black hole (BH) solution in the GR, modified gravity also have characteristic solutions with theory to theory. That is why to find a new solution in the theory of gravitation is important.

The $2+1$ gravity has been suggested by Deser and Jackiw [1]. This theory has the characteristic structure as follows. At first, lower dimensional theory of gravitation has fewer unknown functions than higher dimensional theory. That is why to derive new exact solutions in $2+1$ space-time is more easier than higher dimensional one. Next, Weyl tensor always vanishes in the gravity theory which has only 3 or fewer dimensional space-time. In addition, the lack of degrees of freedom tends to gravitational wave does not propagate. Carlip reviewed such as interesting properties of the $2+1$ gravity [2].

This theory came to attract attention after the discovery of BTZ solution [3]. A solution represents the black hole which is in the AdS space-time. Note that BTZ black hole is not asymptotically flat like Schwarzschild black hole. After their research, a lot of solutions were derived with many types of matter fields in the $2+1$ space-time.

In the present article, let us treat non linear electric field (NLEF) and dilaton field as matters. To the past research, Cataldo find exact solutions with Born-Infeld [4] type and $|\mathcal{F}^{3/4}|$ type NLEF [5] , [6]. Note that $\mathcal{F}$ is the Maxwell invariant which is defined as $\mathcal{F} = F_{\mu\nu}F^{\mu\nu}$ where $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ is the electric tensor in which A_μ is vector potential. Gurtug found an exact solution with $|\mathcal{F}^k|$ (k is a real number) type NLEF which is called Maxwell Power (MP) theory [7]. They are circular symmetric static solution with electric field. Moreover, solutions in the general dimension are derived by some authors [8], [9], [10]. Then, Hendi found that MP type NLEF is related to the spherically symmetric solution of the $F(R) = R^k$ type modified gravity [11]. Recently, new solutions were derived with scalar hair in the AdS space-time [12].

To treat the dilaton field is meaning full because it is related to the open and the closed bosonic string theory (on our paper, we however neglect the Kolb-Ramond antisymmetric tensor field). Dilaton field is also important for the theory of gravitation because it has correspondence to the modified gravity. Since to obtain solutions of the theory is important, researchers find solutions in the theory of gravitation with dilaton field as follows. Gibbons and Maeda found a electrically charged black hole solution in $3+1$ space-time [13]. In the $2+1$ gravity, Sá and Lemos found the black hole solution without electric field [14]. Then, Chan and Mann discovered some solutions with linear electric field [15]. After their research, Yamazaki and Ida introduced Born-Infeld

*u003wa@yamaguchi-u.ac.jp

type NLEF and found a new solution [16]. However, MP type NLEF with dilaton field had not considered. That is why we try to find new solutions of the $2+1$ gravity which is coupled to MP type NLEF with dilaton field.

In this article, we take natural unit which is $c = 8\pi G = 1$, where c is speed of light, G is Newton constant in the $2+1$ space-time. Additionally, we set the metric signature as $(-++)$.

2 Circular symmetric static solutions of the $2+1$ gravity which is coupled to $\mathcal{F}^k$ type NLEF with dilaton field

2.1 Action and basic field equations

The action of the $2+1$ gravity which is coupled to $\mathcal{F}^k$ type NLEF with dilaton field is

$$I = \frac{1}{2}\int d^3x\sqrt{-g} \times \left\{R - \alpha\left(\partial\phi\right)^2 - e^{-\phi}\left|\mathcal{F}\right|^k - 2e^{\beta\phi}\Lambda\right\}. \tag{1}$$

Where g is determinant of the metric $g_{\mu\nu}$, R is Ricci scalar, ϕ is the dilaton field, α and β are constants. We get Einstein equation when we take the variation of $g_{\mu\nu}$,

$$G^\mu_\nu + \Lambda e^{\beta\phi}\delta^\mu_\nu = T^\mu_\nu\,, \tag{2}$$

where T^μ_ν is energy-momentum tensor

$$T^\mu_\nu = \alpha\partial^\mu\phi\partial_\nu\phi - \frac{\alpha}{2}\left(\partial\phi\right)^2\delta^\mu_\nu - \left|\mathcal{F}\right|^k\frac{e^{-\phi}}{2}\left(\delta^\mu_\nu - \frac{4kF^{\mu\nu}F_{\mu\nu}}{\mathcal{F}}\right). \tag{3}$$

On the other hand, let us take the variation of (1) respect to ϕ, we get the following dilaton equation

$$\alpha\nabla_\mu\left(\partial^\mu\phi\right) + \frac{e^{-\phi}}{2}\left|\mathcal{F}\right|^k - \beta e^{\beta\phi}\Lambda = 0\,, \tag{4}$$

In addition, electric field equation is derived from the variation of (1) which is respect to A_μ

$$\partial_\mu\left(\sqrt{-g}e^{-\phi}F^{\mu\nu}\left|\mathcal{F}\right|^{k-1}\right) = 0\,. \tag{5}$$

2.2 Static circular symmetric solutions

Now, let us solve above equations under the circular symmetric static ansatz. We set metric ansatz as

$$ds^2 = -f(r)^2dt^2 + \frac{e^{2\delta(r)}dr^2}{f(r)^2} + r^2d\theta^2\,, \tag{6}$$

where $f(r)$, $\delta(r)$ are unknown functions which is depended only radial coordinate r. Following discussion, we assume dilaton field is depended only r. Beside, electric field ansatz is

$$\mathbf{F} = e^{\phi(r)+\delta(r)}E(r)dt\wedge dr\,, \tag{7}$$

where $E(r)$ and $\phi(r)$ are unkown functions which is depended only r.

Then, Einstein equations will be

$$\frac{e^{-2\delta(r)}\left(f^2\delta_r - ff_r\right)}{r} = \frac{\alpha}{2}e^{-2\delta}f^2\left(\frac{d\phi}{dr}\right)^2 + \frac{1-2k}{2}e^{-\phi}\left|\mathcal{F}\right|^k + e^{\beta\phi}\Lambda\,, \tag{8}$$

$$\frac{e^{-2\delta(r)}ff_r}{r} = \frac{\alpha}{2}e^{-2\delta}f^2\left(\frac{d\phi}{dr}\right)^2 - \frac{1-2k}{2}e^{-\phi}\left|\mathcal{F}\right|^k - e^{\beta\phi}\Lambda\,, \tag{9}$$

$$e^{-2\delta(r)}\left(ff_r\delta_r - ff_{rr} - (f_r)^2\right) = \frac{\alpha}{2}e^{-2\delta}f^2\left(\frac{d\phi}{dr}\right)^2 - \frac{1}{2}e^{-\phi}\left|\mathcal{F}\right|^k + e^{\beta\phi}\Lambda\,. \tag{10}$$

When we subtract (9) from (8), we get

$$\frac{1}{r}\frac{d\delta}{dr} = \alpha\left(\frac{d\phi}{dr}\right)^2. \tag{11}$$

In this paper, we assume function $\delta(r)$ as

$$\delta(r) = n\ln\left(\frac{r}{r_*}\right), \tag{12}$$

which is embraced in article [16]. Here n and r_* are constants. Thus, we can determinant the dilaton field ϕ as

$$\phi(r) = \pm\sqrt{\frac{n}{\alpha}}\ln\left(\frac{r}{r_0}\right), \tag{13}$$

where r_0 is some constant.

Next, let us discuss the electric field equation. We can treat equation (5) analytically, then

$$E^{2k-1}(r) = \frac{q}{r}\left(\frac{r_0}{r}\right)^{\pm2\sqrt{n/\alpha}(k-1)}, \tag{14}$$

where q is an integration constant which is related to the charge of the space-time. The $k=1$ case, when the electric field $E(r)$ proportional to $\ln(r)$, was already discussed by Chan [15].

To determine the relation of constants, we calculate Einstein equations and dilaton field. At first, take the summation (8) and (9), then

$$\frac{e^{-2\delta}(f^2\delta_r - 2ff_r)}{r} = (1-2k)e^{-\phi}\left|\mathcal{F}\right|^k + 2e^{\beta\phi}\Lambda\,. \tag{15}$$

On the other hand, from the dilaton field equation, we get

$$\begin{aligned}&\alpha e^{-2\delta}\left\{\frac{f^2}{r}-f^2\delta_r+2ff_r\right\}\left(\frac{d\phi}{dr}\right)\\&+\alpha e^{-2\delta}f^2\frac{d^2\phi}{dr^2}+\frac{e^{-\phi}}{2}\left|\mathcal{F}\right|^k-\beta e^{\beta\phi}\Lambda=0\,.\end{aligned}\tag{16}$$

Thus we get the following relation when we use equation (12), (13) and (15), (16)

$$\begin{aligned}&\left\{\pm\sqrt{n\alpha}(1-2k)-\frac{1}{2}\right\}e^{-\phi}|\mathcal{F}|^k\\&+\left(\pm2\sqrt{n\alpha}+\beta\right)e^{\beta\phi}\Lambda=0\,.\end{aligned}\tag{17}$$

This equation implies that there are two types of the solution in our discussion. We call the "first class" solution which means the coefficients $e^{-\phi}|\mathcal{F}|^k$ and $e^{\beta\phi}$ vanishes at non zero Λ;

$$-\beta=\pm2\sqrt{n\alpha}\,,\tag{18}$$

$$\beta\,(1-2k)+1=0\,.\tag{19}$$

This class of the solutions are not consistent to $k=1/2$ type MP theory because they are not consistent to the coefficient of $e^{-\phi}|\mathcal{F}|^k$ equal to zero.

Other types of the solution which we call "second class" solutions mean

$$e^{-\phi}|\mathcal{F}|^k=-\frac{\pm2\sqrt{n\alpha}+\beta}{\pm\sqrt{n\alpha}(1-2k)-1/2}\Lambda e^{\beta\phi}\,.\tag{20}$$

This means that coefficients do not vanish. It also means we could get the consistent solution at $k=1/2$ case. Hereafter, we will get the solutions in the explicit form.

2.2.1 The first class solutions

We now consider the first class solutions. In this case, we get the following differential equation

$$\begin{aligned}&\frac{2f}{r}\frac{df}{dr}-n\frac{f^2}{r^2}\\&=(2k-1)\left(\frac{r}{r_*}\right)^{2n}\left(\frac{r}{r_0}\right)^{8nk^2(1-k)/(2k-1)}\\&\times\left(\frac{q}{r}\right)^{2k/(2k-1)}-2\left(\frac{r_0}{r_*}\right)^{2n}\Lambda\,.\end{aligned}\tag{21}$$

This equation can solve analytically, then

$$\begin{aligned}f(r)=&-\left(\frac{r_0}{r_*}\right)^{2n}\frac{2\Lambda r^2}{2-n}-mr^n\\&+\left(\frac{r_0}{r_*}\right)^{2n}\frac{r^2(2k-1)^2}{2(k-1)(1-4nk^2)-n(2k-1)}\\&\times\left(\frac{r}{r_0}\right)^{8nk^2(1-k)/(2k-1)}\left(\frac{q}{r}\right)^{2k/(2k-1)}\,,\end{aligned}\tag{22}$$

where m is an constant. Again, this solution is not consistent to the $k=1/2$ MP theory because of the constrain (19).

2.2.2 The second class solutions

In the second class solution, the electric part $e^{-\phi}|\mathcal{F}|^k$ and cosmological part $e^{\beta\phi}$ have the same r dependence. This condition restricts constants as

$$\pm\sqrt{\frac{n}{\alpha}}=\frac{2k}{1+\beta(1-2k)}\,.\tag{23}$$

Then, equation (20) becomes

$$\begin{aligned}E^{2k}=&-\frac{4k\alpha+\beta(1+\beta(1-2k))}{4k\alpha(1-2k)-\beta(1-2k)-1}2\Lambda\\&\times e^{(\beta-(2k-1))\phi}\,.\end{aligned}\tag{24}$$

After some calculation, we get the following differential equation

$$\begin{aligned}&\frac{2f}{r}\frac{df}{dr}-n\frac{f^2}{r^2}\\&=2\frac{\beta^2(1-2k)^2+8k\alpha(1-2k)-1}{4k\alpha(1-2k)-\beta(1-2k)-1}\Lambda e^{\beta\phi+2\delta}\,.\end{aligned}\tag{25}$$

This equation can solve analytically as

$$\begin{aligned}f^2=&-mr^\alpha-\Lambda r^2\frac{\beta^2(1-2k)^2+8k\alpha(1-2k)-1}{4k\alpha(1-2k)-\beta(1-2k)-1}\\&\times\frac{(1+\beta(1-2k))^2}{2k^2\alpha+(1+\beta(1-2k))(1+\beta(1-k))}\\&\times\left(\frac{r}{r_0}\right)^{2k\beta/(1+\beta(2k-1))}\left(\frac{r}{r_*}\right)^{8k^2\alpha/(1+\beta(1-2k))^2}\,,\end{aligned}\tag{26}$$

where m is an constant.

The solution which belongs to second class is consistent to the $k=1/2$ MP theory. In the $k=1/2$ theory, solutions of the field equations are

$$r_0=q\ ,\ \ \pm\sqrt{\frac{n}{\alpha}}=1\,,\tag{27}$$

$$\phi(r)=\ln\left(\frac{r}{q}\right)\,,\tag{28}$$

$$\delta(r)=\alpha\ln\left(\frac{r}{r_*}\right)\,,\tag{29}$$

$$E(r)=2(2\alpha+\beta)\left(\frac{r}{q}\right)^\beta\Lambda\,,\tag{30}$$

$$f^2(r)=\frac{-2\Lambda r^2}{2+\alpha+\beta}\left(\frac{r}{r_*}\right)^{2\alpha}\left(\frac{r}{q}\right)^\beta-mr^\alpha\,.\tag{31}$$

These solutions are consistent to the solutions of the field equations from the action

$$I = \frac{1}{2}\int d^3x\sqrt{-g} \times \left\{R - \alpha\left(\partial\phi\right)^2 - e^{-\phi}\sqrt{|\mathcal{F}|} - 2e^{\beta\phi}\Lambda\right\}, \tag{32}$$

under the circular symmetric ansatz (6) and (7). It is characteristic structure of the second class solutions. In the past work, special electric field ansatz that treated by Mazharimousavi [17] is also consistent to the $k = 1/2$ type MP theory. However, $k = 1/2$ type MP theory has no consistent solution without dilaton field or first class solutions with circular symmetric ansatz which adopted in our paper.

3 Fundamental structure of the space-time

In this chapter, let us consider the fundamental geometric quantity such as scalar invariant. Addition, we investigate the behavior of the metric function $f(r)$.

As was mentioned above, hereafter we will investigate negative cosmological constant case, then

$$\ell = \frac{1}{\sqrt{-\Lambda}}, \tag{33}$$

where ℓ is cosmological radius. Why we consider negative Λ case? At first, asymptotic behavior of the fundamental black hole solution in the $2+1$ gravity that called BTZ black hole is AdS space-time. Moreover, because of the AdS/CFT correspondence [18], it is believed that AdS space-time itself interesting. So, we hope that it is meaning full to include the negative cosmological constant.

3.1 first class solutions

The solution (22)is constructed by power term of r. Then, we can rewrite (22) as

$$f(r)^2 = Ar^2 - mr^n - B\left(\frac{1}{r}\right)^c, \tag{34}$$

where

$$\begin{aligned} A &= \left(\frac{r_0}{r_*}\right)^{2n}\frac{1}{(2-n)\ell^2}, \\ B &= \frac{(2k-1)^2}{2(k-1)(1-4nk^2)-n(2k-1)}q^{2k/(2k-1)} \\ &\times r_0^{8nk^2(k-1)/(2k-1)}, \\ c &= \frac{2k}{2k-1}\left(1-4nk^2(1-k)\right). \end{aligned} \tag{35}$$

Then, Fundamental geometric quantity such as Ricci scalar of the solution (22) reads to

$$R = -\frac{6Ar^2 + (c^2-5c+6)Br^{2-c} - mn(n+1)r^n}{r^2}. \tag{36}$$

The singular behavior of the Ricci scalar is depended on n and k (equal to α and β) becomes as follows. i) a singular point at origin. ii) a singular point at infinity. iii) two singular points at origin and infinity. Higher order curvature invariant such as$R^{\mu\nu}R_{\mu\nu}$ and $R^{\mu\nu\sigma\lambda}R_{\mu\nu\sigma\lambda}$ also has same structure. Thus, our new solution (22) has physical singular point at $r = 0$ or $r = \infty$ that depends on n and k. In addition, it is possible to create a black hole solution with some fine parameter sets. To get a black hole solution, we will choice i) type physical singular behavior. Then, parameters must satisfy $n - 2 \leq 0$, $C \leq 0$. They are equal to

$$n \leq 2\ ,\ n \leq \frac{1}{4k^2(1-k)}. \tag{37}$$



Figure 1: The behavior of the metric function when singular point lies at $r = 0$. The horizontal line represents radial position r, and vertical line represents the metric function $f(r)$. The parameter sets are $\ell = r_0 = r_* = 1$ and $n = 1$ with $k = 3/4$. Top line, middle line and bottom line means $m = 0.1,\ 0.85,\ 2$ respectively.

The behavior of the metric function in this case is shown in Figure 1 and Figure 2. We cannot derive the horizon radius with analytical method in generally, however, there are horizons at $f(r_H) = 0$ where r_H is its radius at some parameter sets. From Figure 1, we notice that there are two horizons and the space-time is similar to the R-N black hole. Thus, the parameter sets in Figure 1 could generate R-N like black holes. On the other hand, some parameter sets construct the metric function like Figure 2. Then, the space-time is similar to the Schwarzschild-AdS black hole. That is why we conclude that our



Figure 2: The behavior of the metric function when singular point lies at $r = 0$. The parameter sets are $\ell = r_0 = r_* = 1$ and $n = 0.1$ with $k = 3/4$. Left line, middle line and right line means $m = 0.1,\ 2,\ 3$ respectively.

new solutions can generate some different types of black hole.

3.2 second class solutions

Here, we consider about second class solutions as first class solutions. Second class solutions (26) can rewrite as

$$f(r)^2 = -mr^n + Cr^2\left(\frac{1}{r}\right)^d \tag{38}$$

where

$$\begin{aligned} C = &\frac{1}{\ell^2}\frac{\beta^2(1-2k)^2 + 8k\alpha(1-2k) - 1}{4k\alpha(1-2k) - \beta(1-2k) - 1} \\ &\times \frac{(1+\beta(1-2k))^2}{2k^2\alpha + (1+\beta(1-2k))(1+\beta(1-k))} \\ &\times \left(\frac{1}{r_0}\right)^{2\beta k/(1+\beta(2k-1))}\left(\frac{1}{r_*}\right)^{8\alpha k^2/(1+\beta(1-2k))^2}, \end{aligned} \tag{39}$$

$$d = -\frac{2\beta k}{1+\beta(1-2k)} - \frac{8\alpha k^2}{1+\beta(1-2k))^2}\,. \tag{40}$$

Then, Ricci scalar reads to

$$R = -\frac{(d^2-5d+6)Cr^2r^{-d} - mn(n+1)r^n}{r^2}\,. \tag{41}$$

The feature of the Ricci scalar is as follows. i) To diverge at $r = 0$, power terms of r must proportional to $1/r$. This reads to $0 \le d$ and $n - 2 \le 0$. This case our new solution has one physical singular point at origin. ii) To diverge at $r = \infty$, power terms of r has to be proportional to r. This reads $d \le 0$ and $0 \le n - 2$. This case our new solution has one physical singular point at infinity. iii) Both $d = 0$ and $n = 2$, there is no singular point. iv) In other case, Ricci scalar will diverge at $r = 0$ and $r = \infty$. Then, solution has two physical singular point at origin and infinity. Higher order curvature invariant such as $R^{\mu\nu}R_{\mu\nu}$ and $R^{\mu\nu\sigma\lambda}R_{\mu\nu\sigma\lambda}$ also has same structure. Thus, our new solution (26) has physical singular point at $r = 0$ or $r = \infty$ that depends on theory to theory.



Figure 3: The behavior of the metric function when singular point lies at $r = 0$. The parameter sets are $\ell = r_0 = r_* = 1$ and $\alpha = 1/4$, $\beta = 1$ with $k = 2$. Left line, middle line and right line means $m = 4,\ 1,\ 0.5$ respectively.

An example of the behavior of the metric function is shown in Figure 3. We notice that there is one horizon in the figure. In the second class solutions, we can derive the horizon radius as

$$r_H = \left(\frac{m}{C}\right)^{1/(2-d-n)}\,. \tag{42}$$

From the figure 3, we find that the space-time is similar to the black hole in the de-Sitter space-time. That is why we conclude that second class solutions can also generate the black hole.

4 Conclusion and Remarks

We derive new solutions of the $2+1$ space-time that coupled to MP type NLEF with dilaton field. We could derive new circular symmetric static solutions in general k case. There are two types solutions that we called "first class" and "second class". Especially, solutions that belong to second class, we could get consistent solution at $k = 1/2$ theory that has no consistent circular symmetric electric solution without dilaton field. From the singular analyze and behavior of the metric function, we notice that some parameter sets will generate black holes.

The future work that we have to do as follows. At first, we have to investigate more detail structure of the solution. One of them is thermodynamics of the

solution. In the BTZ black hole, thermodynamical consideration has been performed and the behavior of thermodynamic quantities are investigated. The space-time which we derived in our article should be studied along with the similar thermodynamical context. Consider the physical structure such as mechanics of our new solutions seems to interesting. Physical phenomena such as lensing effect is also interesting. Above analyze will derive physical viewpoints of our new solutions.

In addition, to derive more new solutions is important. We treat electric solutions in our article. Then, we did not derive magnetic solutions. So considering the magnetic type solution is meaningful. Second, to consider the rotating effect seems important. It is known that spinning case has more interesting property. That is why we have to treat point symmetric type ansatz in the future work. It seems meaningful to do the same works in higher dimensional case. We have to find solutions of higher dimensional Einstein gravity witch is coupled to non linear electromagnetic field with dilaton field.

Acknowledgments

I would like to thank K. Shiraishi for helpful discussions. I also thank K. Kobayashi for reading the manuscript.

References

[1] S. Deser, R. Jackiw, Annals Phys. **153** (1984) 405-416 .

[2] S. Carlip, Class. Quant. Grav. **12**, 2835 (1995) .

[3] M. Bañados, C. Teitelboim and J. Zanelli, Phys. Rev. Lett. **69**, 1849 (1992) .

[4] M. Born and L. Infeld, Proc. Roy. Soc. Lond. **A144**, 425 -451 (1934) .

[5] M.Cataldo, A. Garcia, Phys.Lett. **B456** (1999) 28-33 .

[6] M. Cataldo, A. Garcia, Phys. Rev. **D61**, 084003 (2000) .

[7] O. Gurtug, S. H. Mazharimousavi, M. Halilsoy, Phys. Rev. **D85**. 104004 (2012) .

[8] M. Hassaine and C. Martinez, Phys. Rev. **D75**, 027502 (2007).

[9] M. Hassaine and C. Martinez, Class. Quant. Grav. **25**, 195023 (2008).

[10] H. Maeda, M. Hassaine and C. Martinez, Phys. Rev. **D79**, 044012 (2009).

[11] S. H. Hendi, Phys. Lett. **B690**, 220 (2010) .

[12] Wei Xu and De-Cheng Zou, "(2+1)-dimensional charged black holes with scalar hair in Einstein-Power-Maxwell Theory", [arXiv:1408.1998v1] .

[13] Gibbons G W and Maeda K, Nucl. Phys. **B298** (1988) 741 .

[14] P. M. Sá and J. P. S. Lemos, Phys. Lett. **B423** (1996) 125-138 .

[15] K. C. K. Chan and R. B. Mann, Phys. Rev. **D50** (1994) 6385-6393 .

[16] R. Yamazaki and D. Ida, Phys. Rev. **D64** 024009 (2001) .

[17] S. H. Mazharimousavi and M. Hailsoy, Ozay Gurtug, Eur. Phys. J. **C74**, 2735 (2014) .

[18] Juan M. Maldacena, Adv. Theor. Math. Phys. **2** (1998) 231-252 .